

A large sample test for the length of memory of stationary symmetric stable random fields via nonsingular $\mathbb{Z}^d$ -actions

Ayan Bhattacharya and Parthanil Roy

Statistics and Mathematics Unit, Indian Statistical Institute

Abstract

Based on the ratio of two block maxima, we propose a large sample test for the length of memory of a stationary symmetric α -stable discrete parameter random field. We show that the power function converges to one as the sample-size increases to infinity under various classes of alternatives having longer memory in the sense of [22]. Ergodic theory of nonsingular $\mathbb{Z}^d$ -actions play a very important role in the design and analysis of our large sample test.

Key words and phrases. Long range dependence, stationary S α S random field, non-singular group action, extreme value theory, statistical hypothesis testing.

2010 Mathematics Subject Classification. Primary 60G10, 60G60, 62H15; Secondary 37A40.

1 Introduction and Preliminaries

A random field $\mathbf{X} = \{X(\mathbf{t}), \mathbf{t} \in \mathbb{Z}^d\}$ is called a stationary, *symmetric α -stable* (S α S) random field if every finite linear combination $\sum_{i=1}^k a_i X_{\mathbf{t}_i + \mathbf{s}}$ is an S α S random variable whose distribution does not depend on $\mathbf{s}$. Here we shall consider the non-Gaussian case (i.e., $0 < \alpha < 2$) unless mentioned otherwise.

Long range dependence is a very important property that has been observed in many real-life processes. By long range dependence of the the random field $\mathbf{X}$, we mean the dependence between the observations $X(\mathbf{t})$ which are far separated in $\mathbf{t}$. This concept was introduced in order to study the measurements of the water flow in Nile river by famous British hydrologist Hurst (see [8] and [9]). In literature, several definitions of long range dependence are available. For a detailed discussion on long range dependence, we refer to [23] and the references therein.

In a seminal work [22], the notion of long memory for a stationary S α S random process has been introduced based on a dichotomy in long run behavior of the partial maxima. A partition of the underlying parameter space (formally defined later) has been suggested

in the aforementioned reference which causes the dichotomy. This dichotomy has been studied for $d \geq 2$ in [21]. Phase transitions in many other probabilistic features of stationary SaaS random fields have been connected to the same partition of the parameter space; see e.g., [14], [16], [20], [6].

The fact that the law of $\mathbf{X}$ is invariant under the group action of shift transformation on the index set $\mathbb{Z}^d$ (stationarity) and certain rigidity properties of L^α spaces ($0 < \alpha < 2$) are used in [18] (for $d = 1$) and [19] (for $d \geq 2$) to show that there always exists an integral representation of the form

$$X(\mathbf{t}) \stackrel{d}{=} \int_{\mathbb{E}} a_{\mathbf{t}}(x) \left(\frac{d m \circ \phi_{\mathbf{t}}}{d m}(x) \right)^{1/\alpha} f \circ \phi_{\mathbf{t}}(x) M(dx), \quad (1.1)$$

where M is an SaaS random measure on a standard Borel space $(\mathbb{E}, \mathcal{E})$ with σ -finite control measure m , $f \in L^\alpha(\mathbb{E}, m)$ (a deterministic function), $\{\phi_{\mathbf{t}}\}$ is a non-singular $\mathbb{Z}^d$ -action on $(\mathbb{E}, m)$ (i.e., each $\phi_{\mathbf{t}} : \mathbb{E} \rightarrow \mathbb{E}$ is measurable and invertible, ϕ_0 is the identity map, $\phi_{\mathbf{t}_1} \circ \phi_{\mathbf{t}_2} = \phi_{\mathbf{t}_1 + \mathbf{t}_2}$ for all $\mathbf{t}_1, \mathbf{t}_2 \in \mathbb{Z}^d$ and each $m \circ \phi_{\mathbf{t}}^{-1}$ is an equivalent measure of m) and $\{c_{\mathbf{t}}\}$ is a measurable cocycle for the nonsingular action $\{\phi_{\mathbf{t}}\}$ taking values in $\{+1, -1\}$ (i.e., each $c_{\mathbf{t}} : \mathbb{E} \rightarrow \{+1, -1\}$ is measurable map such that for all $\mathbf{t}_1, \mathbf{t}_2 \in \mathbb{Z}^d$, $c_{\mathbf{t}_1 + \mathbf{t}_2}(x) = c_{\mathbf{t}_2}(x) c_{\mathbf{t}_1}(\phi_{\mathbf{t}_2}(x))$ for all $x \in \mathbb{E}$).

As a stationary SaaS random field can be uniquely specified in terms of a function in $L^\alpha(\mathbb{E}, m)$, a nonsingular action and a cocycle, we consider the following parameter space for a stationary SaaS random field

$$\Theta = \left\{ (f, \{\phi_{\mathbf{t}}\}, \{c_{\mathbf{t}}\}) : f \in L^\alpha(\mathbb{E}, m), \{\phi_{\mathbf{t}}\} \text{ is a nonsingular action, } \{c_{\mathbf{t}}\} \text{ is a cocycle} \right\}. \quad (1.2)$$

Now based on the nonsingular action, we can get a decomposition of $\mathbb{E}$ (into two subsets) which is known as Hopf decomposition as described below. A set W is called a wandering set for the nonsingular $\mathbb{Z}^d$ -action $\{\phi_t\}$ on $(\mathbb{E}, m)$ if $\{\phi_t(W) : t \in \mathbb{Z}^d\}$ is pairwise disjoint collection of subsets of $\mathbb{E}$. Following Proposition 1.6.1 in [1], we get that $\mathbb{E}$ can be decomposed into two disjoint and invariant (with respect to $\{\phi_t\}$) subsets $\mathcal{C}$ and $\mathcal{D}$ such that for some wandering set $W \subset \mathbb{E}$, $\mathcal{D} = \cup_{t \in \mathbb{Z}^d} \phi_t(W)$ and $\mathcal{C}$ does not have any wandering set of positive measure. $\mathcal{C}$ and $\mathcal{D}$ are called the conservative and dissipative parts of $\{\phi_{\mathbf{t}}\}$, respectively. If $\mathbb{E} = \mathcal{C}$, then we call the nonsingular $\mathbb{Z}^d$ -action $\{\phi_t\}$ conservative. If $\mathbb{E} = \mathcal{D}$, then $\{\phi_t\}$ is called dissipative.

Following [18], [19] and [21], and denoting the integrand in (1.1) by $f_{\mathbf{t}}(x)$,

$$X(\mathbf{t}) \stackrel{d}{=} \int_{\mathcal{C}} f_{\mathbf{t}}(x) M(dx) + \int_{\mathcal{D}} f_{\mathbf{t}}(x) M(dx) =: X^{\mathcal{C}}(\mathbf{t}) + X^{\mathcal{D}}(\mathbf{t}), \quad \mathbf{t} \in \mathbb{Z}^d, \quad (1.3)$$

where $\mathbf{X}^{\mathcal{C}} = \{X^{\mathcal{C}}(\mathbf{t}), \mathbf{t} \in \mathbb{Z}^d\}$ and $\mathbf{X}^{\mathcal{D}} = \{X^{\mathcal{D}}(\mathbf{t}), \mathbf{t} \in \mathbb{Z}^d\}$ are two independent stationary SaaS random field generated by conservative and dissipative nonsingular $\mathbb{Z}^d$ -actions, respectively. It is important to note that the stationary SaaS random field generated by a

dissipative nonsingular $\mathbb{Z}^d$ -action admits mixed moving average representation (see [26] and (1.7) below).

Based on the notion of partial block maxima, it has been established in [22] and [21] that stationary S α S random fields generated by conservative actions have longer memory than those generated by a nonsingular action with a non trivial dissipative part. This has formalized the intuition that “conservative action keeps coming back” (i.e., same value of the random measure M contributes to the observations $X(\mathbf{t})$ which are far separated in $\mathbf{t}$) and hence induces longer memory. Let for all $n \in \mathbb{N}$,

$$\text{Box}(n) = \{\mathbf{j} = (j_1, \dots, j_d) \in \mathbb{Z}^d : |j_i| \leq n \text{ for } 1 \leq i \leq d\} \quad (1.4)$$

be the block containing the origin with size $(2n+1)^d$ in $\mathbb{Z}^d$. We define the partial block maxima for the stationary S α S random field $\mathbf{X}$ as

$$M_n = \max_{\mathbf{j} \in \text{Box}(n)} |X(\mathbf{j})|, \quad n \in \mathbb{N}.$$

The asymptotic behaviour of the partial block maxima M_n is related to the deterministic sequence

$$B_n = \left(\int_{\mathbb{E}} \max_{\mathbf{j} \in \text{Box}(n)} |f_{\mathbf{j}}(x)|^\alpha m(\mathrm{d}x) \right)^{1/\alpha}. \quad (1.5)$$

Note that by Corollary 4.4.6 of [24], B_n is completely specified by the parameters associated to the S α S random field and does not depend on the choice of the integral representation. We shall recall the results on rate of growth of $\{B_n\}$ from [21] (Proposition 4.1). It is expected that the rate of growth of B_n will be slower if the underlying group action is conservative. Indeed, if $\{\phi_{\mathbf{t}} : \mathbf{t} \in \mathbb{Z}^d\}$ is conservative, then

$$\lim_{n \rightarrow \infty} \frac{1}{(2n+1)^{d/\alpha}} B_n = 0. \quad (1.6)$$

In the other case, we need the mixed moving average representation to describe the limit. A stable random field is called a mixed moving average (see [26]) if it is of the form

$$\mathbf{X} \stackrel{d}{=} \left\{ \int_{W \times \mathbb{Z}^d} f(u, \mathbf{s} - \mathbf{t}) M(\mathrm{d}u, \mathrm{d}\mathbf{s}) : \mathbf{t} \in \mathbb{Z}^d \right\}, \quad (1.7)$$

where $f \in L^\alpha(W \times \mathbb{Z}^d, \nu \otimes l)$, l is the counting measure on $\mathbb{Z}^d$, ν is a σ -finite measure on a standard Borel space $(W, \mathcal{W})$ and the control measure m of M equals $\nu \otimes l$. It was shown in [18], [19] and [21] that a stationary S α S random field is generated by a dissipative action if and only if it is a mixed moving average with the integral representation (1.7). In this

case,

$$\lim_{n \rightarrow \infty} \frac{1}{(2n+1)^{d/\alpha}} B_n = \left(\int_W (g(u))^\alpha \nu(du) \right)^{1/\alpha} \in (0, \infty), \quad (1.8)$$

where for every $u \in W$

$$g(u) = \max_{\mathbf{s} \in \mathbb{Z}^d} |f(u, \mathbf{s})|. \quad (1.9)$$

We shall denote the right hand side of (1.8) by $K_{\mathbf{X}}$ which depends solely on $\mathbf{X}$ and not on the integral representation.

Using the above facts, it has been established that, if the SoS random field is not generated by the conservative action then

$$(2n+1)^{-d/\alpha} M_n \Rightarrow C_\alpha^{1/\alpha} K_{\mathbf{X}} Z_\alpha, \quad (1.10)$$

where $K_{\mathbf{X}}$ is as above, Z_α is a standard Fréchet(α) random variable with distribution function

$$\mathbf{P}(Z_\alpha \leq z) = \begin{cases} e^{-z^{-\alpha}} & \text{if } z > 0, \\ 0 & \text{if } z \leq 0, \end{cases}$$

and

$$C_\alpha = \left(\int_0^\infty x^{-\alpha} \sin x \, dx \right) = \begin{cases} \frac{1-\alpha}{\Gamma(2-\alpha) \cos(\pi\alpha/2)} & \text{if } \alpha \neq 1, \\ \frac{2}{\pi} & \text{if } \alpha = 1. \end{cases} \quad (1.11)$$

On the other hand, if the underlying group action is conservative then

$$(2n+1)^{-d/\alpha} M_n \xrightarrow{p} 0. \quad (1.12)$$

See Theorem 4.3 in [21] and Theorem 4.1 in [22].

Note that the dichotomy between (1.10) and (1.12) can be justified by the intuitive reasoning that the longer memory prevents erratic changes in X_t causing the maxima to grow slower. In Gaussian case, this phenomenon occurs in the form of comparison lemma; see, e.g., Corollary 4.2.3 in [13].

The effect of a transition from conservative to dissipative actions has been investigated for various other features of stationary SoS random fields. For example, the ruin probability of negative drifted random walk with steps from a stationary ergodic stable processes, has been studied in [14]. It has been observed that the ruin is more likely if the group action is conservative. The point processes associated to a stationary SoS random field is analysed in [16] (for $d = 1$) and [20] (for $d \geq 2$). It is observed that the point process converges weakly to a Poisson cluster process if the group action is not conservative and in the conservative case, it does not remain tight due to presence of clustering. The large

deviations issues for point process convergence has been addressed in [6], where different large deviation behavior is observed depending on the ergodic theoretic properties of the underlying nonsingular actions.

Stationary S α S random fields have also been studied from statistical perspective (see [24], [11], [10]). Different inference problems associated to the long range dependence for finite and infinite variance processes has been addressed in the literature; see for example, [5], [25], [15], [7], [3], [2], [17] and references therein. There are real-life data such as teletraffic data ([4]) which exhibits heavy-tail phenomenon and long range dependence. Motivated by all these works, the decomposition of the parameter space suggested in [21] and its effect on various probabilistic aspects of S α S random fields, a natural question comes in mind: *is it possible to design a hypothesis testing problem which will detect the presence of long memory in the observed stationary S α S random field?* In the following paragraph, we formulate the problem.

Motivated by [22] and [21] and the other related works mentioned above, we shall consider the following decomposition of the parameter space Θ into Θ_0 and Θ_1 . We define Θ_1 as

$$\Theta_1 = \left\{ \left(f, \{\phi_{\mathbf{t}}\}, \{c_{\mathbf{t}}\} \right) \in \Theta : \{\phi_{\mathbf{t}}\} \text{ is conservative} \right\} \quad (1.13)$$

and $\Theta_0 = \Theta \setminus \Theta_1$. In this article, our aim is to design a large sample statistical test for testing

$$H_0 : \theta \in \Theta_0 \quad \text{against} \quad H_1 : \theta \in \Theta_1 \quad (1.14)$$

where $\theta = (f, \{\phi_{\mathbf{t}}\}, \{c_{\mathbf{t}}\})$ is the parameter associated to the observed stationary S α S random field defined by (1.1).

This article is organized as follows. In Section 2, we shall present a large sample test (based on the ratio of two appropriately scaled block maxima) for testing H_0 vs. H_1 along with its asymptotics under both null and alternative. In particular, our test will become consistent for a reasonably broad class of alternatives. Examples of such alternatives are given in Section 3 along with a numerical experiment. Finally, proofs of our results are discussed in Section 4.

2 Proposed Large Sample Test Based on Block Maxima

Let $\{\mathbf{e}_i : 1 \leq i \leq d\}$ be the d unit vectors in $\mathbb{Z}^d$ such that the i^{th} component of $\mathbf{e}_i$ is 1 and the other components are 0. Let

$$U_n = (2n+1)^{-d/\alpha} \max_{\mathbf{j} \in \text{Box}(n)} |X(\mathbf{j})|$$

and

$$V_n = (2[n^e] + 1)^{-d/\alpha} \max_{\mathbf{j} \in (2n + [n^e])\mathbf{e}_1 + \text{Box}([n^e])} |X(\mathbf{j})|.$$

In other words, U_n is the properly scaled block maxima for $\text{Box}(n)$ containing origin as the centre and V_n is the properly scaled block maxima for shifted $\text{Box}([n^e])$ whose centre is sufficiently separated from $\text{Box}(n)$. To test the hypotheses (1.14), we define the test statistic T_n as the ratio of two partial block maxima U_n and V_n , that is

$$T_n = \frac{U_n}{V_n} = \left(\frac{2[n^e] + 1}{2n + 1} \right)^{d/\alpha} \frac{\max_{\mathbf{j} \in \text{Box}(n)} |X(\mathbf{j})|}{\max_{\mathbf{j} \in (2n + [n^e])\mathbf{e}_1 + \text{Box}([n^e])} |X(\mathbf{j})|}.$$

We shall derive the weak limit of the test statistic T_n under the null hypothesis with the help of following theorem.

Theorem 2.1. *Suppose that the stationary S α S random field $\mathbf{X}$ is generated by a non-conservative action and hence the dissipative part $X^{\mathcal{D}}$ admits a non-trivial moving average representation (1.7), then*

$$(U_n, V_n) \Rightarrow (Y_1, Y_2), \quad (2.1)$$

where Y_i 's are independent copies of Y with distribution function

$$\mathbf{P}(Y \leq y) = \begin{cases} \exp \left\{ -C_\alpha K_{\mathbf{X}}^\alpha y^{-\alpha} \right\} & \text{if } y > 0, \\ 0 & \text{if } y \leq 0. \end{cases} \quad (2.2)$$

and C_α defined in (1.11).

Corollary 2.2. Under the assumptions of Theorem 2.1, $T_n \Rightarrow T$ where T has the distribution function

$$F_T(t) := \mathbf{P}(T \leq t) = \frac{1}{1 + t^{-\alpha}}. \quad (2.3)$$

Proof of Corollary 2.2. Using continuous mapping theorem and the fact that $Y_2 > 0$ almost surely, we get that

$$T_n \Rightarrow T := \frac{Y_1}{Y_2}. \quad (2.4)$$

The distribution of T will be derived using the joint distribution of Y_1 and Y_2 . It is clear that, the joint probability density function is

$$h_{Y_1, Y_2}(y_1, y_2) = (C_\alpha K_{\mathbf{X}}^\alpha)^2 (y_1 y_2)^{-\alpha-1} e^{-C_\alpha K_{\mathbf{X}}^\alpha (y_1^{-\alpha} + y_2^{-\alpha})}, \quad y_1, y_2 > 0.$$

We follow standard substitution procedure by putting $t = y_1 y_2^{-1}$ and $v = y_2$ which in turn gives us $y_1 = tv$ and $y_2 = v$. It is very easy to check that the associated modulus of Jacobian

of transformation is v as $v > 0$. Hence we get the joint distribution of (T, Y_2) as

$$h_{T, Y_2}(t, y_2) = (\alpha C_\alpha K_{\mathbf{X}}^\alpha)^2 t^{-\alpha-1} y_2^{-2\alpha-1} e^{-C_\alpha K_{\mathbf{X}}^\alpha y_2^{-\alpha}(1+t^{-\alpha})}, \quad t > 0, y_2 > 0.$$

Now to get the distribution of T , we have to integrate on the whole range for y_2 . Again using standard substitution

$$z = y_2^{-\alpha}(1+t^{-\alpha})C_\alpha K_{\mathbf{X}}^\alpha$$

we get that

$$\begin{aligned} h_T(t) &= \alpha \frac{t^{-\alpha-1}}{(1+t^{-\alpha})^2} \int_0^\infty z^{2-1} e^{-z} dz \\ &= \frac{\alpha t^{-\alpha-1}}{(1+t^{-\alpha})^2}, \quad t > 0. \end{aligned} \tag{2.5}$$

Hence it is easy to see that (2.3) holds for all $t > 0$. $\square$

We want to compute τ_β such that $\mathbf{P}(T < \tau_\beta) = \beta$. An easy computation yields that,

$$\tau_\beta = \left(\frac{\beta}{1-\beta} \right)^{1/\alpha}. \tag{2.6}$$

The following theorem gives the asymptotics for the test statistic T_n for a very broad class of alternatives.

Theorem 2.3. *Let $\mathbf{X}$ be generated by a conservative $\mathbb{Z}^d$ -action $\{\phi_t\}$. If there exists an increasing sequence of positive real numbers, $\{d_n\}$ such that*

$$d_n = n^{d/\alpha - \eta} L(n), \tag{2.7}$$

where $0 < \eta \leq d/\alpha$ and $L(n)$ is a slowly varying function of n and $\{d_n^{-1} M_n\}_{n \geq 1}$ and $\{d_n M_n^{-1}\}_{n \geq 1}$ are tight sequences of random variables, then we have

$$T_n \xrightarrow{p} 0.$$

So we reject the null hypothesis H_0 against the class of alternatives considered in Theorem 2.3, if $T_n < \tau_\beta$. This gives a large sample level- β test for H_0 against H_1 . Theorem 2.3 ensures that such a test is consistent. In the following section, we shall discuss some examples which satisfy the conditions stated in above theorem. We also derive the empirical power in sub-Gaussian case based on a numerical experiment.

3 Important Classes of Alternatives

In this section, we present a few important examples from the alternative which satisfy the hypotheses of Theorem 2.3 and hence our test becomes consistent.

Example 3.1. We shall first review the basic notions and notations from [21]. Note that the group $R = \{\phi_{\mathbf{t}} : \mathbf{t} \in \mathbb{Z}^d\}$ of invertible non-singular transformations on $(\mathbb{E}, m)$ is a finitely generated abelian group. Define the group homomorphism

$$\Phi : \mathbb{Z}^d \rightarrow R$$

such that $\Phi(\mathbf{t}) = \phi_{\mathbf{t}}$ for all $\mathbf{t} \in \mathbb{Z}^d$. The kernel of this group homomorphism is $\ker(\Phi) = \{\mathbf{t} \in \mathbb{Z}^d : \phi_{\mathbf{t}} = id_{\mathbb{E}}\}$ where $id_{\mathbb{E}}$ denotes the identity map on $\mathbb{E}$. Being a subgroup of $\mathbb{Z}^d$, $\ker(\Phi)$ is a free abelian group. By first isomorphism theorem of groups, we have

$$R \simeq \mathbb{Z}^d / \ker(\Phi).$$

Because of structure theorem for finitely generated abelian groups (Theorem 8.5 in [12]), R can be written as the direct sum of a free abelian group $\bar{F}$ (the free part) and a finite abelian group $\bar{N}$ (the torsion part). So we get

$$R = \bar{F} \oplus \bar{N}.$$

We assume that $1 \leq \text{rank}(\bar{F}) = p < d$. Since $\bar{F}$ is free, there exists an injective group homomorphism

$$\Psi : \bar{F} \rightarrow \mathbb{Z}^d$$

such that $\Phi \circ \Psi = id_{\bar{F}}$. Clearly $F = \Psi(\bar{F})$ is a free subgroup of $\mathbb{Z}^d$ of rank p .

F should be regarded as the effective index set and its rank p becomes the effective dimension of the random field. It was shown in [21] that

$$\frac{1}{(2n+1)^{p/\alpha}} M_n \Rightarrow \begin{cases} C_{\mathbf{X}} Z_{\alpha} & \text{if } \{\phi_{\mathbf{t}}\}_{\mathbf{t} \in F} \text{ is not conservative,} \\ 0 & \text{if } \{\phi_{\mathbf{t}}\}_{\mathbf{t} \in F} \text{ is conservative.} \end{cases}$$

In the above setup, if $1 \leq p < d$, and $\{\phi_{\mathbf{t}}\}_{\mathbf{t} \in \mathbb{Z}^d}$ is not conservative, then using Theorem 2.3 with $d_n = (2n+1)^{p/\alpha}$, we get that

$$T_n \xrightarrow{p} 0$$

as $n \rightarrow \infty$. In particular, the level- β test that rejects when $T_n < \tau_{\beta}$ is consistent.

Example 3.2. We consider a stationary S α S random field indexed by $\mathbb{Z}^2$, with the $\mathbb{Z}^2$ action $\{\phi_{(i,j)}\}_{(i,j) \in \mathbb{Z}^2}$ on $\mathbb{E} = \mathbb{R}$ given by

$$\phi_{(i,j)}(x) = x + i + j\sqrt{2}, \quad x \in \mathbb{R}$$

with m as Lebesgue measure on $\mathbb{R}$. From Example 6.3 in [21], it is clear that

$$\frac{1}{n^{1/\alpha}} M_n \Rightarrow \left((1 + \sqrt{2}) C_{\alpha} \right)^{1/\alpha} Z_{\alpha}.$$

Hence Theorem 2.3 with $d_n = n^{1/\alpha}$ applies and we get

$$T_n \xrightarrow{p} 0$$

as $n \rightarrow \infty$. So the test rejects the null hypothesis H_0 if $T_n < \tau_\beta$ is consistent.

Example 3.3. Consider a random field which has an integral representation of the following form

$$X(\mathbf{j}) = \int_{\mathbb{R}^{\mathbb{Z}^d}} g_{\mathbf{j}} dM, \quad \mathbf{j} \in \mathbb{Z}^d \quad (3.1)$$

where M is an S α S random measure on $\mathbb{R}^{\mathbb{Z}^d}$ whose control measure m is a probability measure under which the projections $g_{\mathbf{j}} : \mathbf{j} \in \mathbb{Z}^d$ are i.i.d. random variables with finite absolute α^{th} moment.

First we consider the case where under m , $\{g_{\mathbf{j}} : \mathbf{j} \in \mathbb{Z}^d\}$ is i.i.d. positive Pareto random variables with

$$m(g_0 > x) = \begin{cases} x^{-\gamma} & \text{if } x \geq 1, \\ 1 & \text{if } x < 1. \end{cases}$$

for some $\gamma > \alpha$. From Example 6.1 in [21], we get that

$$B_n \sim c_{p,\gamma}^{1/\alpha} 2^{d/\gamma} n^{d/\gamma} \text{ as } n \rightarrow \infty$$

for some positive constant $c_{p,\gamma}$ and $B_n^{-1}M_n$ converges weakly to Frechét random variable. So Theorem 2.3 applies with $d_n = n^{d/\gamma}$ and we get

$$T_n \xrightarrow{p} 0$$

as $n \rightarrow \infty$. Hence the level- β test rejects H_0 when $T_n < \tau_\beta$, is consistent.

Now we consider the special case where under m , $\{g_{\mathbf{j}} : \mathbf{j} \in \mathbb{Z}^d\}$ is a sequence of i.i.d. standard normal random variables. Then $\{X_{\mathbf{j}}\}_{\mathbf{j} \in \mathbb{Z}^d}$ has the same distribution as the process $\{c_\alpha A^{1/2} G_{\mathbf{j}}\}_{\mathbf{j} \in \mathbb{Z}^d}$, where $G_{\mathbf{j}}$'s are i.i.d. standard Gaussian random variables, A is a positive $\alpha/2$ -stable random variable independent of $\{G_{\mathbf{j}} : \mathbf{j} \in \mathbb{Z}^d\}$ with Laplace transform $\mathbf{E}(e^{-tA}) = e^{-t^{\alpha/2}}$ and $c_\alpha = \sqrt{2} \left(\mathbf{E}(|G_0|^\alpha) \right)^{1/\alpha}$; see section 3.7 in [24]. Then from Example 6.1 in [21], we get that

$$B_n \sim \sqrt{2d \log 2n}$$

such that

$$B_n^{-1}M_n \Rightarrow A^{1/2},$$

which is a positive random variable.

So we can apply Theorem 2.3 with $d_n = \sqrt{2d \log 2n}$ and obtain

$$T_n \xrightarrow{p} 0$$

as $n \rightarrow \infty$. Hence level- β test that rejects H_0 if $T_n < \tau_\beta$ is consistent .

3.1 Numerical Experiment

In the following simulation, we shall discuss the empirical power of the test for a sub-Gaussian random field indexed by $\mathbb{Z}^2$. We shall see that the empirical power increases as n becomes large. Based on the following tables, we shall suggest a relation between the value of ϱ and α that influences the empirical power.

ϱ	$\alpha = .7$			$\alpha = .9$			$\alpha = 1.1$		
	$n = 80$	$n = 90$	$n = 100$	$n = 80$	$n = 90$	$n = 100$	$n = 80$	$n = 90$	$n = 100$
0.61	1	1	1	1	1	1	1	1	1
0.62	1	1	1	1	1	1	1	1	1
0.63	1	1	1	1	1	1	1	1	1
0.64	1	1	1	1	1	1	1	1	1
0.65	1	1	1	1	1	1	1	1	1
0.66	1	1	1	1	1	1	1	1	1
0.67	1	1	1	1	1	1	0.9975	1	1
0.68	1	1	1	0.9975	1	1	0.9950	1	1
0.69	1	1	1	0.9975	1	1	0.9850	1	0.9975
0.70	1	1	1	0.9875	0.9975	1	0.9600	0.9975	0.9900

Table 1: Empirical power for $\alpha = 0.7, 0.9$ and 1.1 .

ϱ	$\alpha = 1.3$			$\alpha = 1.5$			$\alpha = 1.7$		
	$n = 80$	$n = 90$	$n = 100$	$n = 80$	$n = 90$	$n = 100$	$n = 80$	$n = 90$	$n = 100$
0.61	1	1	1	1	1	1	1	1	1
0.62	1	1	1	1	1	1	1	1	0.9975
0.63	1	1	1	0.9975	1	1	1	1	0.995
0.64	1	1	1	0.9975	1	1	0.9925	1	0.995
0.65	1	1	1	0.9925	1	1	0.9875	0.9925	0.9975
0.66	1	1	1	0.995	1	0.995	0.9650	0.9725	0.9875
0.67	1	1	1	0.9700	0.9825	0.9975	0.9125	0.9825	1
0.68	0.9750	0.9875	0.9875	0.9325	0.9650	0.9925	0.8700	0.9300	0.9575
0.69	0.9500	0.9575	0.9875	0.8825	0.9150	0.9600	0.7625	0.8925	0.9175
0.70	0.8775	0.9375	0.9775	0.7625	0.8650	0.9375	0.6800	0.8125	0.8725

Table 2: Empirical power for $\alpha = 1.3, 1.5$ and 1.7 .

It is clear from the above tables that if we use small values of ϱ , then the rejection will be very frequent and hence power will be very high for each fixed α . On the other hand, a large value of ϱ results in fewer rejection and so power decreases for a fixed α . Observe that the empirical power decreases as α increases for every fixed ϱ . So it seems that we need to choose a smaller value of ϱ as α increases. So there is an inverse relation between ϱ and α . Hence the choice of ϱ is not uniform in α . Note that as n increases, the empirical power increases to 1 for all values of ϱ and α confirming the consistency of the proposed test.

4 Proofs of Theorems 2.1 and 2.3

4.1 Proof of Theorem 2.1

Proof of 2.1. Without loss of generality, we shall assume that $\mathbf{X}$ admits moving average representation. This is because under our hypothesis, we can use the decomposition (1.3) with a non-trivial dissipative part and the conservative part does not contribute to the maxima after scaling. In particular, this means that

$$X(\mathbf{j}) = \int_W \int_{\mathbb{Z}^d} f(u, \mathbf{v} - \mathbf{j}) M(\mathrm{d} u, \mathrm{d} \mathbf{v}), \quad \mathbf{j} \in \mathbb{Z}^d,$$

where M is an S α S random measure on $W \times \mathbb{Z}^d$ with control measure $m = \nu \otimes l$ on $\mathcal{B}(W \times \mathbb{Z}^d)$ where l is counting measure on $\mathbb{Z}^d$. Also $f \in L^\alpha(W \times \mathbb{Z}^d, \nu \otimes l)$. Let $\text{Box}(L) = \{\mathbf{j} \in \mathbb{Z}^d : |j_1| \leq L, \dots, |j_d| \leq L\}$ i.e., it is an L neighbourhood around the origin. Define

$$X(\mathbf{j}, L) = \int_W \int_{\mathbb{Z}^d} f(u, \mathbf{v} - \mathbf{j}) \mathbb{1}_{W \times \text{Box}(L)}(w, \mathbf{v} - \mathbf{j}) M(\mathrm{d} u, \mathrm{d} \mathbf{v}) \quad (4.1)$$

for all positive integer L . Define

$$M_n(L) = \max \left\{ |X(\mathbf{j}, L)| : \mathbf{j} \in \text{Box}(n) \right\} \quad (4.2)$$

$$\text{and} \quad \mathfrak{M}_n(L) = \max \left\{ |X(\mathbf{j}, L)| : \mathbf{j} \in (2n + [n^\varrho])\mathbf{e}_1 + \text{Box}([n^\varrho]) \right\}. \quad (4.3)$$

Fix $L \in \mathbb{N}$. It is important to observe that, as an easy consequence of Theorem 4.3 in [21], we have

$$\frac{1}{(2n+1)^{d/\alpha}} M_n(L) \Rightarrow Y_1(L),$$

where $Y_1(L)$ is a positive random variable with distribution function

$$\mathbf{P}(Y_1(L) \leq y) = \exp \left\{ -C_\alpha K_{\mathbf{X}}^\alpha(L) y^{-\alpha} \right\} \quad (4.4)$$

with

$$K_{\mathbf{X}}^\alpha(L) = \int_W \sup_{\mathbf{j} \in \text{Box}(L)} |f(w, \mathbf{j})|^\alpha \nu(dw).$$

Similar facts lead to the observation that $(2[n^\ell] + 1)^{-d/\alpha} \mathfrak{M}_n$ converges weakly to a random variable with same distribution as that of $Y_1(L)$. It is important to note that for all $n \geq 2L + 1$, we have $\{X(\mathbf{j}, L) : \mathbf{j} \in \text{Box}(n)\}$ and $\{X(\mathbf{j}, L) : \mathbf{j} \in (2n + [n^\ell])\mathbf{e}_1 + \text{Box}([n^\ell])\}$ are independent random vectors which follows from Theorem 3.5.3 in [24]. So M_n and $\mathfrak{M}_n$ are independent for all $n \geq 2L + 1$. Combining these facts we get that

$$\left(\frac{1}{(2n+1)^{d/\alpha}} M_n(L), \frac{1}{(2[n^\ell]+1)^{d/\alpha}} \mathfrak{M}_n(L) \right) \Rightarrow (Y_1(L), Y_2(L))$$

where $Y_1(L)$ and $Y_2(L)$ are independently and identically distributed with law as specified in (4.4). It is easy to see that as $L \rightarrow \infty$, $K_{\mathbf{X}(L)} \rightarrow K_{\mathbf{X}}$. So we have

$$(Y_1(L), Y_2(L)) \Rightarrow (Y_1, Y_2)$$

as $L \rightarrow \infty$.

Now it only remains to show that for every fixed $\epsilon > 0$,

$$\lim_{L \rightarrow \infty} \limsup_{n \rightarrow \infty} \mathbf{P} \left(\frac{1}{(2n+1)^{d/\alpha}} |M_n - M_n(L)| + \frac{1}{(2[n^\ell]+1)^{d/\alpha}} |\mathfrak{M}_n - \mathfrak{M}_n(L)| > \epsilon \right) = 0. \quad (4.5)$$

To show (4.5), it is enough to show that

$$\lim_{L \rightarrow \infty} \limsup_{n \rightarrow \infty} \mathbf{P} \left(\frac{1}{(2n+1)^{d/\alpha}} |M_n - M_n(L)| > \epsilon/2 \right) = 0.$$

Recall that

$$B_n = \left(\int_{\mathbb{E}} \max_{\mathbf{j} \in \text{Box}(n)} |f(w, \mathbf{j})|^\alpha m(dw) \right)^{1/\alpha}$$

and define a new probability measure η_n on $\mathbb{E} = W \times \mathbb{Z}^d$ for every fixed n ,

$$\frac{d\eta_n}{dm}(w, \mathbf{j}) = B_n^{-\alpha} \max_{\mathbf{j} \in \text{Box}(n)} |f(w, \mathbf{j})|^\alpha. \quad (4.6)$$

Using Theorem 3.5.6 and Corollary 3.10.4 from [24], we know that for $\mathbf{j} \in \text{Box}(n)$,

$$X(\mathbf{j}) \stackrel{d}{=} C_\alpha^{1/\alpha} \sum_{i=1}^{\infty} \varepsilon_i \Gamma_i^{-1/\alpha} f(U_i^{(n)}, \mathbf{V}_i^{(n)} - \mathbf{j}), \quad \mathbf{j} \in \text{Box}(n),$$

where C_α is a constant as specified in (1.11), $\{\varepsilon_i : i \geq 1\}$ is a collection of i.i.d. $\{\pm 1\}$ -valued symmetric random variables, $\{\Gamma_i : i \geq 1\}$ is the collection of arrival times of the unit rate Poisson process and $\{(U_i^{(n)}, \mathbf{V}_i^{(n)}) : i \geq 1\}$ is a collection of i.i.d. $\mathbb{E} = W \times \mathbb{Z}^d$ -valued random

variables with common law η_n for every fixed n . It is straight forward to check that

$$X(\mathbf{j}, L) \stackrel{d}{=} C_\alpha^{1/\alpha} \sum_{i=1}^{\infty} \varepsilon_i \Gamma_i^{-1/\alpha} f(U_i^{(n)}, \mathbf{V}_i^{(n)} - \mathbf{j}) \mathbb{1}_{W \times \text{Box}(L)}(U_i^{(n)}, \mathbf{V}_i^{(n)} - \mathbf{j}), \quad \mathbf{j} \in \text{Box}(n).$$

Now note that

$$\begin{aligned} & \max_{\mathbf{j} \in \text{Box}(n)} |X(\mathbf{j})| - \max_{\mathbf{j} \in \text{Box}(n)} |X(\mathbf{j}, L)| \\ &= \max_{\mathbf{j} \in \text{Box}(n)} \left| X(\mathbf{j}, L) \Gamma_i^{-1/\alpha} f(U_i^{(n)}, \mathbf{V}_i^{(n)} - \mathbf{j}) \right. \\ & \quad \left. + C_\alpha^{1/\alpha} \sum_{i=1}^{\infty} \varepsilon_i \mathbb{1}_{W \times (\text{Box}(L))^c}(U_i^{(n)}, \mathbf{V}_i^{(n)} - \mathbf{j}) \right| - \max_{\mathbf{j} \in \text{Box}(n)} |X(\mathbf{j}, L)| \\ &\leq \max_{\mathbf{j} \in \text{Box}(n)} \left| C_\alpha^{1/\alpha} \sum_{i=1}^{\infty} \varepsilon_i \Gamma_i^{-1/\alpha} f(U_i^{(n)}, \mathbf{V}_i^{(n)} - \mathbf{j}) \mathbb{1}_{W \times (\text{Box}(L))^c}(U_i^{(n)}, \mathbf{V}_i^{(n)} - \mathbf{j}) \right| \end{aligned} \quad (4.7)$$

where the last inequality follows from the fact that

$$\max_{\mathbf{j} \in \mathbb{Z}^d} (a_{\mathbf{j}} + b_{\mathbf{j}}) \leq \max_{\mathbf{j} \in \mathbb{Z}^d} a_{\mathbf{j}} + \max_{\mathbf{j} \in \mathbb{Z}^d} b_{\mathbf{j}}$$

for two sequences $\{a_{\mathbf{j}} : \mathbf{j} \in \mathbb{Z}^d\}$ and $\{b_{\mathbf{j}} : \mathbf{j} \in \mathbb{Z}^d\}$ of positive real numbers. Also note that

$$\begin{aligned} & \max_{\mathbf{j} \in \text{Box}(n)} |X(\mathbf{j})| - \max_{\mathbf{j} \in \text{Box}(n)} |X(\mathbf{j}, L)| \\ &= \max_{\mathbf{j} \in \text{Box}(n)} |X(\mathbf{j})| - \max_{\mathbf{j} \in \text{Box}(n)} \left| X(\mathbf{j}) \right. \\ & \quad \left. - C_\alpha^{1/\alpha} \sum_{i=1}^{\infty} \varepsilon_i \Gamma_i^{-1/\alpha} f(U_i^{(n)}, \mathbf{V}_i^{(n)} - \mathbf{j}) \mathbb{1}_{W \times (\text{Box}(L))^c}(U_i^{(n)}, \mathbf{V}_i^{(n)} - \mathbf{j}) \right| \\ &\geq - \max_{\mathbf{j} \in \text{Box}(n)} \left| C_\alpha^{1/\alpha} \sum_{i=1}^{\infty} \varepsilon_i \Gamma_i^{-1/\alpha} f(U_i^{(n)}, \mathbf{V}_i^{(n)} - \mathbf{j}) \mathbb{1}_{W \times (\text{Box}(L))^c}(U_i^{(n)}, \mathbf{V}_i^{(n)} - \mathbf{j}) \right| \end{aligned} \quad (4.8)$$

using the fact that any two sequence of real numbers $\{a_{\mathbf{j}} : \mathbf{j} \in \mathbb{Z}^d\}$ and $\{b_{\mathbf{j}} : \mathbf{j} \in \mathbb{Z}^d\}$ satisfy the following inequality

$$\max_{\mathbf{j} \in \mathbb{Z}^d} |a_{\mathbf{j}}| - \max_{\mathbf{j} \in \mathbb{Z}^d} |a_{\mathbf{j}} - b_{\mathbf{j}}| \geq - \max_{\mathbf{j} \in \mathbb{Z}^d} |b_{\mathbf{j}}|.$$

Now combining the the upper bound in (4.7) and the lower bound obtained in (4.8), we get that

$$\left| \max_{\mathbf{j} \in \text{Box}(n)} |X(\mathbf{j})| - \max_{\mathbf{j} \in \text{Box}(n)} |X(\mathbf{j}, L)| \right| \leq \max_{\mathbf{j} \in \text{Box}(n)} |X^{(c)}(\mathbf{j}, L)|,$$

where

$$X^{(c)}(\mathbf{j}, L) = C_\alpha^{1/\alpha} \sum_{i=1}^{\infty} \varepsilon_i \Gamma_i^{-1/\alpha} f(U_i^{(n)}, \mathbf{V}_i^{(n)} - \mathbf{j}) \mathbb{1}_{W \times (\text{Box}(L))^c}(U_i^{(n)}, \mathbf{V}_i^{(n)} - \mathbf{j}).$$

It is easy to verify that $\{X^{(c)}(\mathbf{j}, L) : \mathbf{j} \in \text{Box}(n)\}$ is a stationary S α S random field which admits mixed moving average representation. Hence we can again use Theorem 4.3 in [21], to get that

$$\frac{1}{(2n+1)^{d/\alpha}} \max_{\mathbf{j} \in \text{Box}(n)} |X^{(c)}(\mathbf{j}, L)| \Rightarrow C_\alpha^{1/\alpha} K_{\mathbf{X}}^{(c)}(L) Z_\alpha,$$

where Z_α is a Frechet random variable with distribution function

$$\mathbf{P}(Z_\alpha < x) = e^{-x^{-\alpha}}$$

and

$$K_{\mathbf{X}}^{(c)}(L) = \int_W \sup_{\mathbf{j} \in \mathbb{Z}^d \setminus \text{Box}(L)} |f(w, \mathbf{j})|^\alpha \nu(dw).$$

Finally we have that

$$\begin{aligned} & \limsup_{n \rightarrow \infty} \mathbf{P} \left((2n+1)^{-d/\alpha} |M_n - M_n(L)| > \epsilon/2 \right) \\ &= \limsup_{n \rightarrow \infty} \mathbf{P} \left((2n+1)^{-d/\alpha} \left| \max_{\mathbf{j} \in \text{Box}(n)} |X(\mathbf{j})| - \max_{\mathbf{j} \in \text{Box}(n)} |X(\mathbf{j}, L)| \right| > \epsilon/2 \right) \\ &\leq \limsup_{n \rightarrow \infty} \mathbf{P} \left((2n+1)^{-d/\alpha} \max_{\mathbf{j} \in \text{Box}(n)} |X^{(c)}(\mathbf{j}, L)| > \epsilon/2 \right) \\ &= \mathbf{P} \left(C_\alpha^{1/\alpha} K_{\mathbf{X}}^{(c)}(L) Z_\alpha > \epsilon/2 \right). \end{aligned} \tag{4.9}$$

It is easy to see that as $L \rightarrow \infty$, $K_{\mathbf{X}}^{(c)}(L) \rightarrow 0$ and hence the expression in (4.9) vanishes. This completes the proof of (4.5). $\square$

4.2 Proof of Theorem 2.3

From the fact that $\{d_n^{-1} M_n\}$ and $\{d_n M_n^{-1}\}$ are tight sequences, it follows using stationarity that $\{(d([n^\varrho]))^{-1} \mathfrak{M}_n\}$ and $\{d([n^\varrho]) \mathfrak{M}_n^{-1}\}$ are tight sequences, where $d(n) := d_n$. Note that as a product of two tight sequences,

$$\frac{d([n^\varrho])}{d_n} \frac{M_n}{\mathfrak{M}_n} \tag{4.10}$$

is also a tight sequence of random variables. Observe that

$$\frac{d([n^\varrho])}{d(n)} \sim n^{(d/\alpha - \eta)(\varrho - 1)} \frac{L([n^\varrho])}{L(n)}$$

as $n \rightarrow \infty$, Note also that

$$T_n = \left(\frac{2[n^\varrho] + 1}{2n + 1} \right)^{d/\alpha} \frac{M_n}{\mathfrak{M}_n}$$

$$\begin{aligned} &\sim n^{d/\alpha(\varrho-1)} \frac{M_n}{\mathfrak{M}_n} \\ &\sim \frac{L(n)}{L([n^\varrho])} n^{\eta(\varrho-1)} \frac{d([n^\varrho])}{d(n)} \frac{M_n}{\mathfrak{M}_n}, \end{aligned}$$

from which the result follows because (4.10) is tight and

$$\frac{L(n)}{L([n^\varrho])} n^{\eta(\varrho-1)} \rightarrow 0$$

using Potter bounds (see, e.g. Resnick(1987)).

Acknowledgements. This research was partially supported by the project RARE-318984 (a Marie Curie FP7 IRSES Fellowship). Parthanil Roy was also supported by Cumulative Professional Development Allowance from Ministry of Human Resource Development, Government of India.

References

- [1] J. Aaronson. *An Introduction to Infinite Ergodic Theory*, volume 50 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, 1997.
- [2] Jan Beran. Maximum likelihood estimation of the differencing parameter for invertible short and long memory autoregressive integrated moving average models. *Journal of the Royal Statistical Society. Series B (Methodological)*, pages 659–672, 1995.
- [3] Jan Beran, RJ Bhansali, and D Ocker. On unified model selection for stationary and nonstationary short-and long-memory autoregressive processes. *Biometrika*, 85(4):921–934, 1998.
- [4] Olivier Cappé, Eric Moulines, J-C Pesquet, Athina P Petropulu, and Xueshi Yang. Long-range dependence and heavy-tail modeling for teletraffic data. *IEEE Signal Processing Magazine*, 19(3):14–27, 2002.
- [5] Pier Luigi Conti, Livia De Giovanni, Stilian A Stoev, and Murad S Taqqu. Confidence intervals for the long memory parameter based on wavelets and resampling. *Statistica Sinica*, pages 559–579, 2008.
- [6] Vicky Fasen and Parthanil Roy. Stable random fields, point processes and large deviations. *Stochastic Processes and their Applications*, 126(3):832–856, 2016.
- [7] Liudas Giraitis and Murad S Taqqu. Whittle estimator for finite-variance non-gaussian time series with long memory. *Annals of Statistics*, pages 178–203, 1999.

- [8] Harold Edwin Hurst. Long-term storage capacity of reservoirs. *Trans. Amer. Soc. Civil Eng.*, 116:770–808, 1951.
- [9] Harold Edwin Hurst. Methods of using long-term storage in reservoirs. *Proceedings of the Institution of Civil Engineers*, 5(5):519–543, 1956.
- [10] Wolfgang Karcher, Elena Shmileva, and Evgeny Spodarev. Extrapolation of stable random fields. *Journal of Multivariate Analysis*, 115:516–536, 2013.
- [11] Wolfgang Karcher and Evgeny Spodarev. Kernel function estimation for stable moving average random fields. 2011.
- [12] Serge Lang. Algebra revised third edition. *Graduate Texts in Mathematics*, 1(211):ALL–ALL, 2002.
- [13] M. R. Leadbetter, Georg Lindgren, and Holger Rootzén. *Extremes and related properties of random sequences and processes*. Springer Series in Statistics. Springer-Verlag, New York-Berlin, 1983.
- [14] Thomas Mikosch and Gennady Samorodnitsky. Ruin probability with claims modeled by a stationary ergodic stable process. *Ann. Probab.*, 28(4):1814–1851, 2000.
- [15] A Montanari, MS Taqqu, and V Teverovsky. Estimating long-range dependence in the presence of periodicity: an empirical study. *Mathematical and computer modelling*, 29(10):217–228, 1999.
- [16] Sidney Resnick and Gennady Samorodnitsky. Point processes associated with stationary stable processes. *Stochastic Process. Appl.*, 114(2):191–209, 2004.
- [17] Peter M Robinson. Log-periodogram regression of time series with long range dependence. *The annals of Statistics*, pages 1048–1072, 1995.
- [18] J. Rosiński. On the structure of stationary stable processes. *The Annals of Probability*, 23:1163–1187, 1995.
- [19] J. Rosiński. Decomposition of stationary α -stable random fields. *Annals of Probability*, 28:1797–1813, 2000.
- [20] Parthanil Roy. Ergodic theory, abelian groups and point processes induced by stable random fields. *The Annals of Probability*, pages 770–793, 2010.
- [21] Parthanil Roy and Gennady Samorodnitsky. Stationary symmetric α -stable discrete parameter random fields. *Journal of Theoretical Probability*, 21(1):212–233, 2008.
- [22] G. Samorodnitsky. Extreme value theory, ergodic theory, and the boundary between short memory and long memory for stationary stable processes. *Annals of Probability*, 32:1438–1468, 2004.

- [23] Gennady Samorodnitsky. Long range dependence. *Foundations and Trends® in Stochastic Systems*, 1(3):163–257, 2007.
- [24] Gennady Samorodnitsky and Murad S. Taqqu. *Stable non-Gaussian random processes*. Stochastic Modeling. Chapman & Hall, New York, 1994. Stochastic models with infinite variance.
- [25] Stilian Stoev and Murad S Taqqu. Wavelet estimation for the hurst parameter in stable processes. In *Processes with long-range correlations*, pages 61–87. Springer, 2003.
- [26] Donatas Surgailis, Jan Rosinski, V Mandrekar, and Stamatis Cambanis. Stable mixed moving averages. *Probability Theory and Related Fields*, 97(4):543–558, 1993.

Ayan Bhattacharya (ayanbhattacharya.isi@gmail.com)

Parthanil Roy (parthanil.roy@gmail.com)

Statistics and Mathematics Unit

Indian Statistical Institute

203 B. T. Road

Kolkata 700108, India.